



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The circadian clock that controls gene expression in Arabidopsis is tissue specific

Citation for published version:

Thain, SC, Murtas, G, Lynn, JR, McGrath, RB & Millar, AJ 2002, 'The circadian clock that controls gene expression in Arabidopsis is tissue specific', *Plant physiology*, vol. 130, no. 1, pp. 102-10.
<https://doi.org/10.1104/pp.005405>

Digital Object Identifier (DOI):

[10.1104/pp.005405](https://doi.org/10.1104/pp.005405)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Plant physiology

Publisher Rights Statement:

RoMEO green

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The Circadian Clock That Controls Gene Expression in Arabidopsis Is Tissue Specific¹

Simon C. Thain², Giovanni Murtas³, James R. Lynn, Robert. B. McGrath, and Andrew J. Millar*

Department of Biological Sciences, University of Warwick, Coventry CV4 7AL, United Kingdom (S.C.T., G.M., A.J.M.); Horticulture Research International, Wellesbourne, Warwick CV35 9EF, United Kingdom (J.R.L.); and Laboratory of Molecular Biology, The Rockefeller University, 1230 York Avenue, New York, New York 10021 (R.B.M.)

The expression of *CHALCONE SYNTHASE* (*CHS*) expression is an important control step in the biosynthesis of flavonoids, which are major photoprotectants in plants. *CHS* transcription is regulated by endogenous programs and in response to environmental signals. Luciferase reporter gene fusions showed that the *CHS* promoter is controlled by the circadian clock both in roots and in aerial organs of transgenic Arabidopsis plants. The period of rhythmic *CHS* expression differs from the previously described rhythm of chlorophyll *a/b*-binding protein (*CAB*) gene expression, indicating that *CHS* is controlled by a distinct circadian clock. The difference in period is maintained in the wild-type Arabidopsis accessions tested and in the *de-etiolated 1* and *timing of CAB expression 1* mutants. These clock-affecting mutations alter the rhythms of both *CAB* and *CHS* markers, indicating that a similar (if not identical) circadian clock mechanism controls these rhythms. The distinct tissue distribution of *CAB* and *CHS* expression suggests that the properties of the circadian clock differ among plant tissues. Several animal organs also exhibit heterogeneous circadian properties in culture but are believed to be synchronized in vivo. The fact that differing periods are manifest in intact plants supports our proposal that spatially separated copies of the plant circadian clock are at most weakly coupled, if not functionally independent. This autonomy has apparently permitted tissue-specific specialization of circadian timing.

Light is a key environmental signal for plants, regulating gene expression and development (Neff et al., 2000). Changes in fluence rate and light quality can occur unpredictably and rapidly during the day but have an underlying day-night cycle. Plants have evolved a circadian timing system that allows the anticipation of this predictable rhythm. When plants are deprived of environmental time cues and placed in constant ("free running") environmental conditions, circadian rhythms persist with a period of around 24 h, often for many days (Millar, 1999; McClung, 2000; Murtas and Millar, 2000; Johnson, 2001). Within the circadian system of the whole organism, the term "circadian oscillator" has been used to denote the parts of the system responsible for rhythm generation. Light-dark signals entrain the oscillator via input phototransduction pathways, synchroniz-

ing its phase with the environmental light-dark cycle and also affecting its period. Rhythmic output from the oscillator controls a large number of physiological processes in plants (Lumsden and Millar, 1998). The abundance of 2% to 6% of RNA transcripts in Arabidopsis plants was scored as circadian-regulated in two recent microarray analyses, for example (Harmer et al., 2000; Schaffer et al., 2001).

The rhythmic expression of chlorophyll *a/b*-binding protein (*CAB* or *Light-Harvesting Complex [LHCB]*) genes has often been used as a marker for circadian regulation in plants (for review, see Fejes and Nagy, 1998), especially using firefly (*Photinus pyralis*) luciferase (*LUC*) reporter fusions (Millar et al., 1995a). *CAB* genes are strongly expressed in the mesophyll cells of photosynthetically active organs and in epidermal guard cells. However, physiological analysis shows that the plant circadian system comprises many copies of the circadian clock, with at least one clock in all the major plant organs and possibly one in most cells (Gorton et al., 1989; Kim et al., 1993; Mayer and Fischer, 1994). We use the term "circadian clock" to denote the smallest complete timing unit (comprising an oscillator with light input and output to overt rhythms). *CAB* rhythms, thus, reveal only a subset of clocks in the plant circadian system. The distributed circadian clocks are thought to function autonomously, because many circadian rhythms persist in isolated tissue explants (for example, Vaadia, 1960; Simon et al., 1976; Engelmann and Johnsson, 1978; Thain et al., 2000). Within a single

¹ This work was supported by the Biotechnology and Biological Science Research Council (graduate studentship to S.C.T. and grant no. BI11209 to A.J.M.). The Warwick imaging facility is supported by the Gatsby Charitable Foundation, the Royal Society, and the Biotechnology and Biological Science Research Council (grants to A.J.M.).

² Present address: Division of Environmental and Applied Biology, School of Life Sciences, University of Dundee, Dundee DD1 4HN, Scotland, UK.

³ Present address: Biogemma, Napoli, Italy.

* Corresponding author; e-mail Andrew.Millar@warwick.ac.uk; fax 44-24-7652-3701.

Article, publication date, and citation information can be found at www.plantphysiol.org/cgi/doi/10.1104/pp.005405.

plant, we have shown that various organs can maintain rhythmic expression of the same gene set to different phases (Thain et al., 2000). Central circadian pacemakers and communicated rhythmic signals, therefore, have little influence on plant rhythms, at least for the gene expression rhythms tested (Thain et al., 2000). This is in contrast to their well-documented involvement in mammalian rhythms (van Esseveldt et al., 2000; Yamazaki et al., 2000).

The autonomy of plant clocks implies that, if the timing properties of the circadian clocks varied among tissues, those differences would be reflected in each tissue's circadian rhythms. Such specialization in timing might be advantageous (Roenneberg and Mittag, 1996). It is unclear how much circadian timing actually varies among plant tissues and what are the molecular causes of such variation. Differences in circadian period between plant rhythms have been reported (Hennessey and Field, 1992; Millar et al., 1995a; Fowler et al., 1999; Sai and Johnson, 1999). The rhythms in question were not only expressed in different cells but also had overtly unrelated mechanisms (leaf movement and *CAB* gene expression, for example).

To investigate the heterogeneity of plant circadian clocks, we required rhythmic markers that can be tested for period under constant conditions, in broader spatial domains than the *CAB* genes. Here, we characterize the rhythmic expression of a *CHS* promoter:reporter gene fusion (*CHS:LUC*). *CHS* is expressed predominantly in epidermal cells of aerial organs and in roots, in Arabidopsis (Chory and Peto, 1990; Kaiser and Batschauer, 1995) as in other species (Schmelzer et al., 1988; Ehmann et al., 1991; Haussuhl et al., 1996). We show that its circadian rhythm has a significantly different period than rhythmic *CAB* expression, implying that *CHS* is controlled by a different circadian clock. To test whether the same molecular components are required for circadian control of *CHS* and *CAB*, we assayed the two markers in mutant backgrounds that are known to alter *CAB* rhythms by different mechanisms. The mutations had very similar effects upon both *CHS* and *CAB* rhythms, indicating that the circadian clocks share similar molecular components. The heterogeneity among circadian rhythms of plant gene expression is likely attributable to tissue-specific modifiers of a common biochemical oscillator, which is present in many if not all Arabidopsis cells.

RESULTS

Expression of *CHS:LUC* in Roots and Leaves

Fusions of firefly *LUC* to the promoters of the white mustard (*Sinapis alba*) chalcone synthase (*mCHS*) and Arabidopsis chalcone synthase (*CHS*) genes were transformed into Arabidopsis plants of the Landsberg *erecta* (*Ler*) and C24 strains, respectively. Figure

1B shows that luminescence driven by the *CHS* promoter was evident in the leaves, hypocotyl, and roots of 12-d-old seedlings. Particularly high expression occurred in the shoot apical region and in young lateral roots. Younger, 5-d-old seedlings showed no *CHS* expression in the hypocotyl (data not shown), consistent with previous reports (Kaiser and Batschauer, 1995; Kaiser et al., 1995). The *mCHS* promoter had an identical pattern of expression (data not shown). In contrast, *CAB:LUC* activity was largely confined to green tissues, with a gradient down the hypocotyl and no detectable activity in the roots (Fig. 1D).

The spatial pattern of *CHS* promoter activity was examined in greater detail in leaf sections from *CHS:LUC* plants and plants carrying a *CHS-GUS* fusion transgene. The *CHS:LUC* signal was highest in the upper epidermis, but both reporters showed patchy expression also in the palisade mesophyll layer (Fig.




Figure 1. Tissue-specific expression of the *CHS* promoter. Reflected-light (A and C) and luminescence (B and D) images of *CHS:LUC* (A and B) and *CAB:LUC* (C and D) seedlings after 12 d of growth in LD (12, 12) of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$. Luminescence video images were processed in false color (see "Materials and Methods"): White and red shades represent the highest photon counts, and darker blues, the lowest. A reflected-light image of the plant is shown to the left of each false color image. E through H, Leaf sections were prepared from plants grown in LD (12, 12) of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 7 d and then transferred to $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 5 d. E, Microtome section of a *CHS*: β -glucuronidase (*GUS*) leaf, stained for *GUS* activity and counter-stained with ruthenium red. F through H, Thick hand section of a *CHS:LUC* leaf: bright field (F), luminescence (G), and overlaid (H) images. E and F, Scale bars represent $60 \mu\text{m}$. E and H, Arrowheads indicate areas where *CHS* expression is very weak or absent from the mesophyll.




Figure 2. Rhythmic luminescence in roots and aerial organs of *Arabidopsis* seedlings expressing *CHS:LUC*. *CHS:Luc* (A, B, C, and E) or *mCHS:LUC* (D) expression was assayed by video imaging (see "Materials and Methods") at the times indicated. A, B, D, and E, Seedlings were grown on vertical agar plates, in 7 d of LD (12, 12) at $150 \mu\text{mol m}^{-2} \text{s}^{-1}$. Seedlings in A and D were transferred (at 0 h) to continuous light of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; seedlings in B were transferred to continuous light of $250 \mu\text{mol m}^{-2} \text{s}^{-1}$; and seedlings in E were transferred first to $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 12 h and then to constant darkness. Seedlings in C were both grown and tested at $250 \mu\text{mol m}^{-2} \text{s}^{-1}$. The data are representative of at least five independent experiments. White box on time axis, Light interval; black box, dark interval. Lv, Luminescence from aerial organs; Rt, luminescence from roots.

1, E and G). The epidermal expression of *CHS:LUC* was about 2-fold higher than the mesophyll expression (in Fig. 1G, for example, average counts per pixel per minute were 0.36 for epidermis and 0.19 for upper mesophyll). A minor contribution from rhythmic *CHS* expression in the mesophyll would not be distinguished in our assays. Luminescence from the mesophyll is appreciably reduced by passage through the epidermis (Wood et al., 2001), indicating that the *CHS:LUC* rhythm preferentially reflects epidermal luminescence. *CHS-GUS* expression was also evident in all cell layers of root cross sections (data not shown).

The *CHS* Expression Rhythm Is Distinct from the *CAB* Expression Rhythm

The strong *CHS:LUC* activity in the root might make this a useful marker for circadian rhythms specifically in this organ, where root pressure and ion fluxes are the only previously reported, rhythmic markers (Vaadia, 1960; Parsons and Kramer, 1974; Gorr et al., 1995; Henzler et al., 1999). The circadian rhythm of *CHS:LUC* activity was tested under several fluence rates of constant light, because *CHS* mRNA levels increase in response to higher fluence rates (Feinbaum and Ausubel, 1988; Peter et al., 1991; Fu-

glevand et al., 1996). Transgenic seedlings were grown for 12 d on vertical agar plates under white light of 60, 150, or 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ fluence rate. Seedlings grown and assayed in white light of 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ showed low expression of *CHS:LUC* and *mCHS:LUC* (data not shown). Luminescence rhythms were measured after the plants were transferred to constant light of either the same fluence rates (Fig. 2, A and C), or of an increased fluence rate (from 150 to 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$; Fig. 2B). *CHS:LUC* activity showed circadian rhythmicity in aerial organs and roots under all conditions. The peak of activity on the 1st d in constant light occurred before predicted dawn under all conditions (at zeitgeber time [ZT] 20–22; ZT is defined as the number of hours since lights-on), 6 or 7 h before the peak of *CAB* expression. Transgenic plants carrying the *mCHS:LUC* fusion in the *Ler* background showed very similar luminescence rhythms (Fig. 2D).

The lighting conditions affected the amplitude of rhythmic *CHS* expression. The high-amplitude rhythm in the 1st d of constant light was followed by a reduction (damping) in the amplitude of the *CHS:LUC* circadian rhythm within two to three cycles. *mCHS:LUC* was less affected (Fig. 2D), possibly because of a difference between the C24 and *Ler* genetic backgrounds. *CAB:LUC* activity continued to cycle with high amplitude under constant light of all fluence rates. In plants transferred from lower to higher fluence rates of light (Fig. 2B), *CAB:LUC* luminescence levels were reduced compared with *CHS:LUC*. When 12-d-old plants expressing *CHS:LUC* were transferred to continuous darkness, transcription from the *CHS* promoter was greatly attenuated (Fig. 2E). A single, high-amplitude peak at the expected phase was followed by rapid damping of *CHS:LUC* activity, with a complete loss of rhythmic amplitude by 72h.

The *CAB* and *CHS* rhythms had different free-running periods under constant light. The lag (phase angle) between the rhythms, therefore, changed progressively during our experiments, most obviously

when *Ler* plants were imaged on the 3rd to 7th d under constant light (Fig. 2D). The altered period was clear: The first peak of *mCHS* expression shown (at 68 h) occurred 10 h before the *CAB* peak (at 78 h), but by 147 h, the *mCHS* peak occurred with or slightly after that for *CAB*. Period estimates were derived from luminescence rhythms measured in the aerial tissues, where both *CHS:LUC* and *CAB:LUC* are expressed (Table I). The period of the *CHS* rhythm (25.4 h) was significantly longer than the period of the *CAB* rhythm (23.7 h). A similar period difference was observed between *CHS* and *CAB* expression rhythms in the *Ler* background (as in Fig. 2D). In contrast, the period of rhythmic *CHS* expression in the root was not significantly different from that in the leaf, in either genetic background (Fig. 2; Table II; data not shown). These results suggested that the circadian system that controlled *CHS* in aerial organs was distinct from that controlling *CAB*.

de-etiolated 1 (det1) and timing of CAB expression 1 (toc1) Mutations Shorten the Period of Both CHS and CAB Rhythms

The *det1* mutant has a severe short-period phenotype for *CAB* expression (Millar et al., 1995b), along with other phenotypes principally related to light signaling (Chory and Peto, 1990; Pepper et al., 1994). *toc1* is a short-period mutant that affects only clock-regulated processes (Millar et al., 1995a; Kreps and Simon, 1997; Somers et al., 1998b; Dowson-Day and Millar, 1999); *TOC1* encodes one of the best candidates for a plant circadian oscillator component (Strayer et al., 2000; Alabadi et al., 2001). The *CHS:LUC* construct was crossed into these mutant backgrounds to test whether these genes are involved in the circadian systems that control both the *CAB* and *CHS* markers. *det1* and *toc1* mutant seedlings expressed *CHS:LUC* rhythmically in both leaves and roots under constant light (Fig. 3A; data not shown). Light-grown *det1* seedlings express the *CHS* promoter in all leaf cell layers, and express *CAB* genes

Table I. Circadian period of gene expression in the aerial organs of *det1*

Plants were maintained as described in the legend to Figure 2B: grown for 7 d under LD (12,12) of 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and transferred to constant light of 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for rhythm assays. Period estimation and statistical comparisons were performed as described in "Materials and Methods." SE are based on the analysis of seven experiments with 129 period estimates (119 degrees of freedom). Genetic backgrounds: C24, Columbia (Col); C24 \times Columbia F2 (C24/Col). The significance levels of *t* tests comparing the mean periods were as follows: ^a*CHS* (+) versus *CAB* (+), $P < 0.0001$. ^b*CHS* (C24) versus *CHS* (C24/Col), $P > 0.4$. ^c*CHS* (+) versus *CHS* (*det1*), $P < 0.0001$. ^d*CHS* (*det1*) versus *CAB* (*det1*), $P < 0.001$.

<i>LUC</i> Marker	Genotype	Background	Period	SE	<i>n</i>
			<i>h</i>		
<i>CAB</i>	+	C24	23.7 ^a	0.36	40
<i>CHS</i>	+	C24	25.4 ^b	0.38	40
<i>CHS</i>	+	C24/Col	25.0 ^c	0.41	20
<i>CHS</i>	<i>det1-1</i>	C24/Col	21.8 ^d	0.48	15
<i>CAB</i>	<i>det1-1</i>	C24/Col	20.7	0.39	14

Table II. Circadian period of *LUC* activity rhythms in *toc1*

Plants were maintained as described for Figure 2B and Table I. Period estimation and statistical comparisons were performed as described in "Materials and Methods." All comparisons of *toc1* versus wild type for a given marker and tissue, $P < 0.0001$; *CHS* in aerial tissues versus roots, $P > 0.15$ in both wild type and *toc1*. All lines are in the C24 background. The significance levels of *t* tests comparing the mean periods were as follows: ^a*CHS* (+) versus *CAB* (+), $P = 0.09$. ^b*CAB* (*toc1*) versus *CHS* (*toc1*) in aerial organs, $P < 0.005$.

<i>LUC</i> Marker	Genotype	Organs	Period	SE	<i>n</i>
			<i>h</i>		
<i>CAB</i>	+	Aerial	24.3 ^a	0.08	4
<i>CHS</i>	+	Aerial	25.1	0.37	5
<i>CAB</i>	<i>toc1-1</i>	Aerial	19.2 ^b	0.25	6
<i>CHS</i>	<i>toc1-1</i>	Aerial	20.6	0.23	5
<i>CHS</i>	+	Root	24.8	0.21	5
<i>CHS</i>	<i>toc1-1</i>	Root	21.2	0.26	4

inappropriately in roots and in aerial tissue (Chory and Peto, 1990). *CAB:LUC* activity in *det1* roots was barely detectable and was too low for us to measure circadian rhythms (data not shown). The *det1* mutation prevents the damping of rhythmic *CAB* expression in darkness (Fig. 3B; Millar et al., 1995b). It had

little effect on *CHS* expression, which damped out in the dark similarly in the mutant (Fig. 3B) and wild type (Fig. 2E). Consistent with previous data, the *toc1* mutation did not affect the mean expression level or damping of *CAB* and *CHS* expression rhythms (data not shown; Millar et al., 1995a).

Quantitative comparisons showed that the period of *CHS* expression was longer than the period of *CAB* expression in both mutant backgrounds (Tables I and II), reinforcing the conclusion that separate clocks control these two promoters. However, both rhythmic markers were very similarly affected by the mutations (Tables I and II), with much shorter periods in mutant progeny than in wild-type controls. This result indicates that *TOC1* and *DET1* function similarly in the circadian clocks that regulate *CAB* and *CHS*.




Figure 3. Circadian rhythms of *CHS:LUC* activity in *det1*. Expression was assayed as in Figure 2. Seedlings were grown in LD (8, 16) at $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ to confirm the *det1* phenotype. Seedlings were transferred to LD (12, 12) at $150 \mu\text{mol m}^{-2} \text{s}^{-1}$, 2 d before the start of the experiment and to constant light of $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ at 0 h (A) or to $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ at 0 h and constant darkness at 12 h (B). Luminescence data were analyzed from whole seedlings, without separate analysis of roots and aerial tissues. The data are representative of at least four independent experiments. White box on time axis, Light interval; black box, dark interval.

DISCUSSION

Chalcone synthase is one of the key biosynthetic enzymes controlling anthocyanin formation. These flavonoid pigments function to protect plant cells from UV radiation and from pathogen attack, act as insect repellents, and are involved in plant-microbe and pollen-pistil signaling (Hahlbrock and Scheel, 1989). We used the noninvasive *LUC* reporter gene in fusions with the *CHS* promoter (*CHS:LUC*), to monitor the dynamic pattern of *CHS* expression (Fig. 1). The rhythmic luminescence of *CHS:LUC* seedlings showed that the circadian clock regulates *CHS* expression at the level of transcription (Fig. 2). *CHS:LUC* represents the first noninvasive molecular marker for circadian rhythms in root tissue.

Regulation of *CHS*

The expression of *CHS* exhibited a very similar circadian rhythm in all tissues (Fig. 2). The phase of the circadian rhythm of *CHS* was earlier than *CAB*, with a peak occurring in the late subjective night (ZT 20–22). Chalcone synthase enzyme activity and mRNA abundance have previously been reported to exhibit diurnal and circadian regulation (Peter et al., 1991; Deikman and Hammer, 1995; Harmer et al.,

2000; Schaffer et al., 2001). The reported phase of peak enzyme activity lags slightly behind the peaks of mRNA abundance and of transcriptional activity, suggesting that the circadian system principally regulates *CHS* expression at the transcriptional level. It may be advantageous for the plant to accumulate photoprotective pigments in advance of the daily photoperiod (Harmer et al., 2000); the timing of *CHS* transcription before dawn is consistent with this notion.

Increasing the fluence rate of white light from 150 to 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ had little effect on the phase or period of *CHS* or *CAB* expression (Fig. 2; data not shown). However, mean expression levels were differentially affected by the lighting conditions. Plants entrained at 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ but assayed at 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ showed reduced levels of *CAB:LUC* activity compared with *CHS:LUC* activity (Fig. 2B), perhaps reflecting a requirement for increased photoprotection and reduced light-harvesting capacity. High-fluence rate light was required to maintain high expression levels of *CHS* transcription in wild-type plants: *CHS:LUC* activity was very low in plants assayed at 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (data not shown) or in darkness (Fig. 2E). The *det1* mutation increases *CAB* expression levels in dark-adapted plants but had little effect on the level of *CHS* expression (Fig. 3B; Chory and Peto, 1990; Millar et al., 1995b). This presumably reflects a differential involvement of *DET1* in the phototransduction pathways that regulate these promoters (Mustilli et al., 1999; Jenkins et al., 2001).

Differences in the Circadian Regulation of *CHS* and *CAB*

The rhythm of *CHS* expression was very similar in roots and in aerial organs under constant light, indicating that the clocks controlling *CHS* in these organs do not differ significantly. The period of *CHS* expression in aerial tissues was approximately 1.5 h longer than the period of *CAB* expression. This difference was maintained in two wild-type accessions and in the *det1* and *toc1* mutants (Tables I and II). Distinct circadian clocks, therefore, control the rhythms of *CAB* and *CHS* expression, although both are nuclear genes that could in principle respond to the same regulator. The phase of a single rhythm (free calcium concentration) has recently been shown to vary among tissues of transgenic tobacco (*Nicotiana tabacum*; Wood et al., 2001). The authors point out that a phase difference could result from tissue-specific clocks or from tissue-specific responses to a single, common clock. Where period differences are observed, the latter interpretation can be ruled out (Sai and Johnson, 1999).

It is not surprising that the period difference is small. Studies on cyanobacteria indicate that a clock with a period that matches the environmental light-

dark cycle provides a competitive advantage (Ouyang et al., 1998). Balancing selection is, therefore, likely to maintain periods in a narrow range around 24 h. Consistent with this notion, we have previously shown that the similar periods of several Arabidopsis accessions are the result of balancing long- and short-period alleles at multiple loci (Swarup et al., 1999).

Period differences under constant conditions are relatively easy to measure. Under light-dark cycles, however, clocks with different periods will entrain to different phases (for example, Ouyang et al., 1998; Somers et al., 1998b). A clock with a longer period (such as that controlling *CHS*) will be set to a later phase, all else being equal. The longer period moves the peak of *CHS* expression away from midnight and toward dawn. If *CAB* was controlled by the same, longer-period clock, then the peak of *CAB* expression would also be later in the day, all else being equal. The delay between the peak of *CHS* expression and the peak of *CAB* expression would thus be greater than we observe. Independent clocks allow the temporal sequence of metabolic processes to be fine-tuned: A smaller delay between the peaks of *CHS* and *CAB* expression might be one example of this. Such a flexible timing system has potential selective advantages (Roenneberg and Mittag, 1996), although these remain to be demonstrated experimentally.

Differential Regulation of Circadian Period

The circadian clocks that control *CAB* and *CHS* might differ fundamentally in their oscillator mechanism, or they might alternatively be separate copies of a common biochemical mechanism with only minor modifications leading to the period difference (Millar, 1998). *det1* and *toc1* mutations are thought to affect the circadian rhythm of *CAB* expression via the input pathway and the oscillator, respectively (Millar et al., 1995a, 1995b; Strayer et al., 2000; Alabadi et al., 2001). Each mutation shortens the circadian period of *CHS* in aerial organs in parallel with *CAB* (Tables I and II). This result shows that both circadian clocks share at least the *TOC1* and *DET1* functions, so the clocks are not radically different. A parsimonious explanation is that the *CHS* and *CAB* promoters are controlled by separate copies of the same clock mechanism (Fig. 4). The characteristics of circadian timing also vary among rodent organs, outside the brain (Yamazaki et al., 2000). Differences in input pathways might be particularly important in that case (Damiola et al., 2000; Stokkan et al., 2001). The same canonical clock genes seem to be involved in diverse anatomical locations (Ripperger et al., 2000).

CHS is expressed principally in the epidermal cell layer of wild-type aerial organs, whereas *CAB* is expressed in the mesophyll layers. It is most likely that their different circadian periods reflect tissue-specific modifications of the clock in epidermal and mesophyll cells, respectively. In support of this conclusion, we




Figure 4. Autonomy and specialization of the circadian clock. A simplified model of the circadian clock is shown in the mesophyll and in the epidermis, with the output markers and clock-related genes tested in this work. The clock is depicted as containing the same components in each case; *TOC1* is thought to function in the oscillator, and *DET1* is thought to affect the light input pathway from the photoreceptors. Rhythmic output from the oscillator is shown regulating transcription factors at the *CAB* or *CHS* promoter. Two hypothetical tissue-specific factors are shown: modification of the light input pathway and an unknown tissue-specific factor (X). One or both modify the function of an oscillator component, resulting in a longer period in the epidermis.

have recently shown that the *PHYTOCHROME B* gene, which is expressed in the epidermis, also has a longer period than *CAB* (A. Hall, L. Kozma-Bognar, F. Nagy, and A.J. Millar, unpublished data). These data imply that epidermal circadian clocks are not tightly coupled to clocks in the mesophyll of the same leaf. We have previously demonstrated that several autonomous clocks exist within the mesophyll of a single leaf (Thain et al., 2000). Taken together, our results suggest that circadian control is local to one or a few cells, not widely coupled within or between tissues, at least for gene expression rhythms in the leaf.

The molecular cause of the observed period difference is unclear. Any circadian input pathway might contribute (Fig. 4). Photoreceptor genes are differentially expressed in the epidermal and mesophyll cell layers (Somers and Quail, 1995), and photoreceptors are known to alter circadian period in a dose-dependent manner (Somers et al., 1998b; Devlin and Kay, 2000). The light input pathways might thus control circadian period in a tissue-specific fashion. Observed distinctions between circadian rhythms suggest that further specialization of the circadian clock might be present in the cells that sense or respond to photoperiodic signals (for example, Salisbury and Denney, 1971; Fowler et al., 1999) or control rhythmic leaf movement (Hennessey and Field, 1992; Fowler et al., 1999; Park et al., 1999). Rhythmic reporters that peak at various phases in restricted spatial patterns are required to determine how much the circadian clock is modulated for specialized timing functions. Recent microarray experiments suggest many candidate promoters (Harmer et al., 2000; Schaffer et al., 2001), which can now be tested in detail using *LUC* fusions.

MATERIALS AND METHODS

Plant Material, Media, and Growth Conditions

Experiments were performed with transgenic *Arabidopsis* seedlings carrying the native firefly (*Photinus pyralis*) *LUC* gene under the control of one of two chalcone synthase promoters. *Arabidopsis* plants of the C24 accession carrying the native *Arabidopsis CHS* promoter fused to the *LUC* gene have been described (Michelet and Chua, 1996). The Columbia transgenic line carrying a fusion of the same *Arabidopsis CHS* promoter fragment to *GUS* was also described previously (Hartmann et al., 1998). Transgenic plants of the *Ler* accession carried the promoter of white mustard (*Sinapis alba*) *CHS1* from -907 to +26 bp (Batschauer et al., 1991), fused to the *LUC* gene in the pMON 721 binary vector backbone. Four independently transformed lines were tested with very similar results (data not shown). The *CAB:LUC* transgene introgressed from C24 into the *Ler* background has been described (Somers et al., 1998b). *CHS:LUC* from the C24 accession was crossed with the *det1-1* mutant in the Columbia background (from Dr. Joanne Chory, Salk Institute). *det1-1* mutant and wild-type plants were selected from the F₂ population on the basis of their morphology; mutant scoring was confirmed in the F₃ generation. The line carrying *CAB:LUC* in the *det1-1* background has been described (Millar et al., 1995b). *CHS:LUC* in C24 was crossed to the *toc1-1* mutant that carries *CAB:LUC* in the C24 background (Millar et al., 1995a). Homozygous mutant F₂ progeny were selected by scoring short-period *CAB:LUC* luminescence in leaves; among these, seedlings carrying *CHS:LUC* were selected by scoring luminescence in roots. The *CAB:LUC* reporter was removed by segregation in the F₃, resulting in greatly reduced luminescence in leaves. The presence of *CHS:LUC* and absence of *CAB:LUC* was confirmed by PCR using transgene-specific primers designed from published sequences (data not shown).

Seedlings were grown at 22°C in 12:12-h light-dark cycles (unless otherwise indicated) on agar medium containing 3% (w/v) Suc under cool-white fluorescent lights of the fluence rates indicated for each figure. Our standard conditions (approximately 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$) gave low *CHS* expression (data not shown); higher fluence rates of 150 to 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ increased expression from these constructs, so all subsequent experiments were conducted within this range.

Localization of *CHS* Expression

Histochemical localization of *GUS* (Fig. 1E) was performed as described (Jefferson et al., 1987). Ruthenium red counterstaining was carried out by standard techniques. Tissue sections were prepared for light microscopy using a Technovit 7100 embedding kit (Kulzer, Wehrheim, Germany). High-resolution imaging of *LUC* activity was performed essentially as described (Hall et al., 2001). The image in Figure 1, G and H, was captured through a Fluor 20x objective (Zeiss, Jena, Germany) by a 30-min exposure on a back-thinned CCD in a liquid-nitrogen cooled camera (LN/CCD-512-TKB with ST133 controller, Roper Scientific Ltd., Marlow, UK). Camera background has been removed from Figure 1G. The specific signal in this image ranges from 2 (dark blue) to 24 (white) counts per pixel. Dark current was undetectable and readout noise was 1.8 counts (sd). The lowest luminescence levels have been clipped in Figure 1H to reveal the tissue outline.

Imaging of *LUC* Bioluminescence Rhythms and Statistical Analysis

After luciferin pretreatment (Millar et al., 1992), seedlings were sprayed before each image with 1 mM D-luciferin (Promega, Madison, WI) in 0.01% (w/v) Triton X-100 and left in darkness for 5 min to allow chlorophyll chemiluminescence to decay. Bioluminescence was detected by ultra-low-light cameras (VIM C2400-47, Hamamatsu, Bridgewater, NJ; and Roper Scientific LN/CCD-512-TKB with ST138 controller), as described (Millar et al., 1992; Michelet and Chua, 1996; Hall et al., 2001). The data from both cameras were similar, although absolute luminescence counts are not directly comparable because of the different image acquisition methods (for example, in Figs. 2 and 3). Data are expressed as detected photons per plant per 25-min image, derived from data from groups of 10 to 20 plants. Data in Figures 2 and 3 have been processed with a three-point, boxcar filter. Period estimates from the raw data were produced by the fast Fourier transform-nonlinear least squares method (FFT-NLLS, Plautz et al., 1997). The first 24 h

of each time course was excluded from the period estimation to exclude any transient effects of the transfer to continuous light.

To derive mean period estimates from the results of seven experiments, the data of Table I were analyzed using residual maximum likelihood (REML) (Patterson and Thompson, 1971) in the statistical package Genstat 5 (Payne et al., 1993). REML can be thought of as a generalization of analysis of variance to unbalanced designs. Data were weighted for analysis by the reciprocal of the estimated variance of the circadian period for the trace, which was derived from FFT-NLLS (Millar et al., 1995a). The data were analyzed with each line taken as a fixed effect and experiment and trace within experiment as random effects. The significance of differences between pairs of treatments were assessed using *t* tests, based on SE of the differences derived from REML, rather than the SE of individual means presented in Table I (Patterson and Thompson, 1971). The mean periods and SE of Table II are derived from the same variance-weighting procedure on period estimates from FFT-NLLS for replicated samples in a separate experiment.

Waveforms in the data that are unusually close to a cosine wave can give very low estimated SEs from FFT-NLLS and, thus, gain disproportionate weight in the variance-weighted means. The analysis was repeated with revised weights that were derived by adding 0.1 to the original, estimated SEs to reduce the effects of such rare estimates. The conclusions of the analysis were not altered by this procedure, and the data presented are weighted using the original estimates.

ACKNOWLEDGMENTS

We thank Profs. Gareth Jenkins and Nam-Hai Chua for helpful discussions and materials and members of the chronobiology group at Warwick for discussions and assistance with imaging experiments.

Received March 12, 2002; accepted April 22, 2002.

LITERATURE CITED

- Alabadi D, Oyama T, Yanovsky MJ, Harmon FG, Mas P, Kay SA (2001) Reciprocal regulation between *TOC1* and *LHY/CCA1*. *Science* **293**: 880–883
- Batschauer A, Ehmann B, Schafer E (1991) Cloning and characterization of a chalcone synthase gene from mustard and its light-dependent expression. *Plant Mol Biol* **16**: 175–185
- Chory J, Peto CA (1990) Mutations in the *DET1* gene affect cell-type-specific expression of light-regulated genes and chloroplast development in *Arabidopsis*. *Proc Natl Acad Sci USA* **87**: 8776–8780
- Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U (2000) Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* **14**: 2950–2961
- Deikman J, Hammer PE (1995) Induction of anthocyanin accumulation by cytokinins in *Arabidopsis thaliana*. *Plant Physiol* **108**: 47–57
- Devlin PF, Kay SA (2000) Cryptochromes are required for phytochrome signaling to the circadian clock but not for rhythmicity. *Plant Cell* **12**: 2499–2509
- Dowson-Day MJ, Millar AJ (1999) Circadian dysfunction causes aberrant hypocotyl elongation patterns in *Arabidopsis*. *Plant J* **17**: 63–71
- Ehmann B, Ocker B, Schafer E (1991) Development-dependent and light-dependent regulation of the expression of two different chalcone synthase transcripts in mustard cotyledons. *Planta* **183**: 416–422
- Engelmann W, Johnsson A (1978) Attenuation of the petal movement rhythm in *Kalanchoe* with light pulses. *Physiol Plant* **43**: 68–76
- Feinbaum RL, Ausubel F (1988) Transcriptional regulation of the *Arabidopsis thaliana* chalcone synthase gene. *Mol Cell Biol* **8**: 1985–1992
- Fejes E, Nagy F (1998) Molecular analysis of circadian clock-regulated gene expression in plants: features of the “output” pathways. In PJ Lumsden, AJ Millar, eds, *Biological Rhythms and Photoperiodism in Plants*. BIOS Scientific, Oxford, pp 99–118
- Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Coupland G, Putterill J (1999) *GIGANTEA*: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J* **18**: 4679–4688
- Fuglevand G, Jackson JA, Jenkins GI (1996) UV-B, UV-A and blue light signal transduction pathways interact synergistically to regulate chalcone synthase expression in *Arabidopsis*. *Plant Cell* **8**: 2347–2357
- Gorr G, Obst G, Doring O, Bottger M (1995) Light-dependent proton excretion of wheat (*Triticum aestivum* L) and maize (*Zea mays* L) roots. *Bot Acta* **108**: 351–357
- Gorton HL, Williams WE, Binns ME, Gemmell CN, Leheny EA, Shepherd AC (1989) Circadian stomatal rhythms in epidermal peels from *Vicia faba*. *Plant Physiol* **90**: 1329–1334
- Hahlbrock K, Scheel D (1989) Physiology and molecular biology of phenylpropanoid metabolism. *Annu Rev Plant Physiol Plant Mol Biol* **40**: 347–369
- Hall A, Kozma-Bognar L, Toth R, Nagy F, Millar AJ (2001) Conditional circadian regulation of *PHYTOCHROME A* gene expression. *Plant Physiol* **127**: 1808–1818
- Harmer SL, Hogenesch JB, Straume M, Chang HS, Han B, Zhu T, Wang X, Kreps JA, Kay SA (2000) Orchestrated transcription of key pathways in *Arabidopsis* by the circadian clock. *Science* **290**: 2110–2113
- Hartmann U, Valentine WJ, Christie JM, Hays J, Jenkins GI, Weisshaar B (1998) Identification of UV/blue light-response elements in the *Arabidopsis thaliana* chalcone synthase promoter using a homologous protoplast transient expression system. *Plant Mol Biol* **36**: 741–754
- Hausuhl K, Rohde W, Weissenböck G (1996) Expression of chalcone synthase genes in coleoptiles and primary leaves of *Secale cereale* L after induction by UV radiation: evidence for a UV-protective role of the coleoptile. *Bot Acta* **109**: 229–238
- Hennessey TL, Field CB (1992) Evidence of multiple circadian oscillators in bean plants. *J Biol Rhythms* **7**: 105–113
- Henzler T, Waterhouse RN, Smyth AJ, Carvajal M, Cooke DT, Schaffner AR, Steudle E, Clarkson DT (1999) Diurnal variations in hydraulic conductivity and root pressure can be correlated with the expression of putative aquaporins in the roots of *Lotus japonicus*. *Planta* **210**: 50–60
- Jefferson RA, Kavanagh TA, Bevan MW (1987) GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J* **6**: 3901–3907
- Jenkins GI, Long JC, Wade HK, Shenton MR, Bibikova TN (2001) UV and blue light signalling: pathways regulating chalcone synthase gene expression in *Arabidopsis*. *New Phytol* **151**: 121–131
- Johnson CH (2001) Endogenous timekeepers in photosynthetic organisms. *Annu Rev Physiol* **63**: 695–728
- Kaiser T, Batschauer A (1995) *cis*-Acting elements of the *CHS1* gene from white mustard controlling promoter activity and spatial patterns of expression. *Plant Mol Biol* **28**: 231–243
- Kaiser T, Emmler K, Kretsch T, Weisshaar B, Schafer E, Batschauer A (1995) Promoter elements of the mustard *CHS1* gene are sufficient for light regulation in transgenic plants. *Plant Mol Biol* **28**: 219–229
- Kim HY, Coté GG, Crain RC (1993) Potassium channels in *Samanea saman* protoplasts controlled by phytochrome and the biological clock. *Science* **260**: 960–962
- Kreps JA, Simon AE (1997) Environmental and genetic effects on circadian clock-regulated gene expression in *Arabidopsis*. *Plant Cell* **9**: 297–304
- Lumsden PJ, Millar AJ, eds (1998) *Biological rhythms and photoperiodism in plants*. BIOS Scientific, Oxford
- Mayer WE, Fischer C (1994) Protoplasts from *Phaseolus coccineus* L. pulvinar motor cells show circadian volume oscillations. *Chronobiol Int* **11**: 156–164
- McClung CR (2000) Circadian rhythms in plants: a millennial view. *Physiol Plant* **109**: 359–371
- Michelet B, Chua NH (1996) Improvement of *Arabidopsis* mutant screens based on luciferase imaging in *planta*. *Plant Mol Biol Rep* **14**: 320–329
- Millar AJ (1998) The cellular organization of circadian rhythms in plants: not one but many clocks. In PJ Lumsden, AJ Millar, eds, *Biological Rhythms and Photoperiodism in Plants*. BIOS Scientific, Oxford, pp 51–68
- Millar AJ (1999) Tansley review no. 103: biological clocks in *Arabidopsis thaliana*. *New Phytol* **141**: 175–197
- Millar AJ, Carré IA, Strayer CA, Chua NH, Kay SA (1995a) Circadian clock mutants in *Arabidopsis* identified by luciferase imaging. *Science* **267**: 1161–1163
- Millar AJ, Short SR, Hiratsuka K, Chua N-H, Kay SA (1992) Firefly luciferase as a reporter of regulated gene expression in higher plants. *Plant Mol Biol Rep* **10**: 324–337

- Millar AJ, Straume M, Chory J, Chua N-H, Kay SA (1995b) The regulation of circadian period by phototransduction pathways in *Arabidopsis*. *Science* **267**: 1163–1166
- Murtas G, Millar AJ (2000) How plants tell the time. *Curr Opin Plant Biol* **3**: 43–46
- Mustilli AC, Fenzi F, Ciliento R, Alfano F, Bowler C (1999) Phenotype of the tomato high pigment-2 mutant is caused by a mutation in the tomato homolog of DEETIOLATED1. *Plant Cell* **11**: 145–157
- Neff MM, Fankhauser C, Chory J (2000) Light: an indicator of time and place. *Genes Dev* **14**: 257–271
- Ouyang Y, Andersson CR, Kondo T, Golden SS, Johnson CH (1998) Resonating circadian clocks enhance fitness in cyanobacteria. *Proc Natl Acad Sci USA* **95**: 8660–8664
- Park DH, Somers DE, Kim YS, Choy YH, Lim HK, Soh MS, Kim HJ, Kay SA, Nam HG (1999) Control of circadian rhythms and photoperiodic flowering by the *Arabidopsis* GIGANTEA gene. *Science* **285**: 1579–1582
- Parsons L, Kramer PJ (1974) Diurnal cycling in root resistance to water movement. *Physiol Plant* **30**: 19–23
- Patterson HD, Thompson R (1971) Recovery of inter-block information when block sizes are unequal. *Biometrika* **58**: 545–554
- Payne RW, Lane P, Digby P, Harding S, Leech P, Morgan G, Todd A, Thompson R, Tunnicliffe W, Welham S et al. (1993) *Genstat 5 Release 3 Reference Manual*. Oxford University Press, Oxford
- Pepper A, Delaney T, Washburn T, Poole D, Chory J (1994) *DET1*, a negative regulator of light-mediated development and gene expression in *Arabidopsis*, encodes a novel nuclear-localized protein. *Cell* **78**: 109–116
- Peter H-J, Kroger-Alef C, Knogge W, Brinkmann K, Weissenböck G (1991) Diurnal periodicity of chalcone-synthase activity during the development of oat primary leaves. *Planta* **183**: 409–415
- Plautz JD, Straume M, Stanewsky R, Jamison CF, Brandes C, Dowse HB, Hall JC, Kay SA (1997) Quantitative analysis of *Drosophila period* gene transcription in living animals. *J Biol Rhythms* **12**: 204–217
- Ripperger JA, Shearman LP, Reppert SM, Schibler U (2000) *CLOCK*, an essential pacemaker component, controls expression of the circadian transcription factor DBP. *Genes Dev* **14**: 679–689
- Roenneberg T, Mittag M (1996) The circadian program of algae. *Semin Cell Dev Biol* **7**: 753–763
- Sai J, Johnson CH (1999) Different circadian oscillators control Ca²⁺ fluxes and *Lhcb* gene expression. *Proc Natl Acad Sci USA* **96**: 11659–11663
- Salisbury FB, Denney A (1971) Separate clocks for leaf movements and photoperiodic flowering in *Xanthium strumarium* L. In M Menaker, ed, *Biochronometry*. National Academy of Sciences, Washington, DC, pp 292–311
- Schaffer R, Landgraf J, Monica A, Simon B, Larson M, Wisman E (2001) Microarray analysis of diurnal and circadian-regulated genes in *Arabidopsis*. *Plant Cell* **13**: 113–123
- Schmelzer E, Jahnen W, Hahlbrock K (1988) In situ localization of light-induced chalcone synthase messenger-RNA, chalcone synthase, and flavonoid end products in epidermal cells of parsley leaves. *Proc Natl Acad Sci USA* **85**: 2989–2993
- Simon E, Satter RL, Galston AW (1976) Circadian rhythmicity in excised *Samanea pulvini*: II. Resetting the clock by phytochrome conversion. *Plant Physiol* **58**: 421–425
- Somers DE, Devlin PF, Kay SA (1998a) Phytochromes and cryptochromes in the entrainment of the *Arabidopsis* circadian clock. *Science* **282**: 1488–1490
- Somers DE, Quail PH (1995) Temporal and spatial expression patterns of *PHYA* and *PHYB* genes in *Arabidopsis*. *Plant J* **7**: 413–427
- Somers DE, Webb AAR, Pearson M, Kay SA (1998b) The short-period mutant, *toc1-1*, alters circadian clock regulation of multiple outputs throughout development in *Arabidopsis thaliana*. *Development* **125**: 485–494
- Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M (2001) Entrainment of the circadian clock in the liver by feeding. *Science* **291**: 490–493
- Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Mas P, Panda S, Kreps JA, Kay SA (2000) Cloning of the *Arabidopsis* clock gene *TOC1*, an autoregulatory response regulator homolog. *Science* **289**: 768–771
- Swarup K, Alonso-Blanco C, Lynn JR, Michaels SD, Amasino RM, Koornneef M, Millar AJ (1999) Natural allelic variation identifies new genes in the *Arabidopsis* circadian system. *Plant J* **20**: 67–77
- Thain SC, Hall A, Millar AJ (2000) Functional independence of circadian clocks that regulate plant gene expression. *Curr Biol* **10**: 951–956
- Vaadia Y (1960) Autonomic diurnal fluctuations in rate of exudation and root pressure of decapitated sunflower plants. *Physiol Plant* **13**: 701–717
- van Esseveldt LKE, Lehman MN, Boer GJ (2000) The suprachiasmatic nucleus and the circadian time-keeping system revisited. *Brain Res Rev* **33**: 34–77
- Wood NT, Haley A, Viry-Moussaid M, Johnson CH, van der Luit AH, Trewavas AJ (2001) The calcium rhythms of different cell types oscillate with different circadian phases. *Plant Physiol* **125**: 787–796
- Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M, Tei H (2000) Resetting central and peripheral circadian oscillators in transgenic rats. *Science* **288**: 682–685