



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Effects of Acute Insulin-Induced Hypoglycemia on Indices of Inflammation Putative mechanism for aggravating vascular disease in diabetes

Citation for published version:

Wright, RJ, Newby, DE, Stirling, D, Ludlam, CA, Macdonald, IA & Frier, BM 2010, 'Effects of Acute Insulin-Induced Hypoglycemia on Indices of Inflammation Putative mechanism for aggravating vascular disease in diabetes', *Diabetes Care*, vol. 33, no. 7, pp. 1591-1597. <https://doi.org/10.2337/dc10-0013>

Digital Object Identifier (DOI):

[10.2337/dc10-0013](https://doi.org/10.2337/dc10-0013)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Diabetes Care

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Effects of Acute Insulin-Induced Hypoglycemia on Indices of Inflammation

Putative mechanism for aggravating vascular disease in diabetes

ROHANA J. WRIGHT, MRCP¹
DAVID E. NEWBY, PHD²
DAVID STIRLING, PHD³

CHRISTOPHER A. LUDLAM, PHD³
IAN A. MACDONALD, PHD⁴
BRIAN M. FRIER, MD¹

OBJECTIVE — To examine the effects of acute insulin-induced hypoglycemia on inflammation, endothelial dysfunction, and platelet activation in adults with and without type 1 diabetes.

RESEARCH DESIGN AND METHODS — We studied 16 nondiabetic adults and 16 subjects with type 1 diabetes during euglycemia (blood glucose 4.5 mmol/l) and hypoglycemia (blood glucose 2.5 mmol/l). Markers of inflammation, thrombosis, and endothelial dysfunction (soluble P-selectin, interleukin-6, von Willebrand factor [vWF], tissue plasminogen activator [tPA], high-sensitivity C-reactive protein [hsCRP], and soluble CD40 ligand [sCD40L]) were measured; platelet-monocyte aggregation and CD40 expression on monocytes were determined using flow cytometry.

RESULTS — In nondiabetic participants, platelet activation occurred after hypoglycemia, with increments in platelet-monocyte aggregation and P-selectin ($P \leq 0.02$). Inflammation was triggered with CD40 expression increasing maximally at 24 h ($3.13 \pm 2.3\%$ vs. $2.06 \pm 1.0\%$) after hypoglycemia ($P = 0.009$). Both sCD40L and hsCRP ($P = 0.02$) increased with a nonsignificant rise in vWF and tPA, indicating a possible endothelial effect. A reduction in sCD40L, tPA, and P-selectin occurred during euglycemia ($P = 0.03$, $P \leq 0.006$, and $P = 0.006$, respectively). In type 1 diabetes, both CD40 expression ($5.54 \pm 4.4\%$ vs. $3.65 \pm 1.8\%$; $P = 0.006$) and plasma sCD40L concentrations increased during hypoglycemia (peak 3.41 ± 3.2 vs. 2.85 ± 2.8 ng/ml; $P = 0.03$). Platelet-monocyte aggregation also increased significantly at 24 h after hypoglycemia ($P = 0.03$). A decline in vWF and P-selectin occurred during euglycemia ($P \leq 0.04$).

CONCLUSIONS — Acute hypoglycemia may provoke upregulation and release of vasoactive substances in adults with and without type 1 diabetes. This may be a putative mechanism for hypoglycemia-induced vascular injury.

Diabetes Care 33:1591–1597, 2010

In people with type 1 diabetes the rapid institution of strict glycemic control aggravates microvascular complications, particularly retinopathy (1). Although attributed to reduced capillary blood flow causing localized ischemia (1), greater exposure to hypoglycemia may have worsened microangiopathy through its putative effects on local vasculature (2). In addition, cardiovascular stress associated with hypoglycemia may precipitate acute macrovascular events in a diseased circulation. While supported by anecdotal reports (3), the increase in cardiovascular mortality in people with type 2 diabetes in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial (4) (and possibly in the Veterans Affairs Diabetes Trial [5]), in which intensive treatment had tripled the frequency of severe hypoglycemia, has caused concern.

tate acute macrovascular events in a diseased circulation. While supported by anecdotal reports (3), the increase in cardiovascular mortality in people with type 2 diabetes in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial (4) (and possibly in the Veterans Affairs Diabetes Trial [5]), in which intensive treatment had tripled the frequency of severe hypoglycemia, has caused concern.

Possible mechanisms by which hypoglycemia may damage blood vessels include changes in regional blood flow, mobilization and activation of neutrophils, platelet activation, and enhanced coagulation and viscosity of the blood (3,6–8). Plasma concentrations of C-reactive protein, interleukin-6 (IL-6), and endothelin-1 increase during hypoglycemia (9–11) and may promote vascular disease (12).

Investigation of processes operating at a cellular level to cause atherosclerosis has focused on the potential influences of vascular inflammation, endothelial dysfunction, coagulation, and platelet activation. The present study sought to determine the effects of acute insulin-induced hypoglycemia on inflammation, coagulation, and platelet and monocyte function in adults with and without type 1 diabetes.

RESEARCH DESIGN AND METHODS

Participants in the study included 16 nondiabetic adult volunteers with no medical history and 16 healthy adults with type 1 diabetes (Table 1). Those with diabetes had no history of hypertension or macrovascular disease, and microvascular disease was excluded. Screening for retinopathy used digital retinal photography, absence of neuropathy was confirmed by clinical examination, and nephropathy was excluded by the absence of microalbuminuria. Subjects with a history of impaired awareness of hypoglycemia or a previous serious reaction to hypoglycemia were excluded. None had a history of head injury, seizure, blackouts, alcohol or drug abuse and psychiatric illness, and their only other medication was the contraceptive pill. Diabetes Control and Complications Trial-aligned A1C was measured using high performance liquid chromatography (nondiabetic reference range 5.0–6.05%; Bio-Rad Laboratories, Munich, Germany); the mean \pm SD of the participants with diabetes was $7.91 \pm 0.92\%$. All gave written informed consent before participation, and the study was approved by the Local Medical Research Ethics Committee.

From the ¹Department of Diabetes, Royal Infirmary of Edinburgh, Edinburgh, U.K.; the ²Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, U.K.; the ³Department of Haematology, Royal Infirmary of Edinburgh, Edinburgh, U.K.; and the ⁴School of Biomedical Sciences, University of Nottingham, Nottingham, U.K.

Corresponding author: Brian M. Frier, brian.frier@luht.scot.nhs.uk.

Received 4 January 2010 and accepted 22 March 2010. DOI: 10.2337/dc10-0013.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

See accompanying original article, p. 1529, and editorial, p. 1686.

Table 1—Baseline demographic characteristics

	Nondiabetic subjects	Subjects with diabetes
<i>n</i>	16	16
Age (years)	28 (26.7–35)	28 (25–37.5)
BMI (kg/m ²)	22.86 ± 2.4	26.40 ± 4.0
Male/female	6/10	7/9
Duration of diabetes (years)	N/A	10 (4.2–19)
A1C (%)	N/A	7.91 ± 0.9

Data are median (interquartile range) and means ± SD unless otherwise indicated.

A modified hyperinsulinemic glucose clamp (13) was used to maintain blood glucose at a predetermined level: euglycemia at 4.5 mmol/l and hypoglycemia at 2.5 mmol/l. Each subject underwent two laboratory sessions, separated by at least 2 weeks (mean 7.2 weeks), of a euglycemic study and a hypoglycemic study in a randomized, counterbalanced fashion.

The participants with type 1 diabetes monitored blood glucose intensively during the 48 h preceding each study, which was postponed if any blood glucose value was <3.5 mmol/l or if symptoms suggestive of hypoglycemia were experienced. After fasting overnight, morning insulin was withheld. A retrograde-intravenous cannula for blood-glucose sampling was inserted into the nondominant hand, which was heated to arterialize the venous blood (14). A cannula in the nondominant antecubital fossa was used to infuse 20% dextrose and soluble insulin (Human Actrapid; Novo Nordisk, Crawley, U.K.) at a constant rate of 1.5 mU/kg/min using a Gemini PCI pump (Alaris Medical Systems, San Diego, CA). The dextrose was infused at a variable rate depending on arterialized blood glucose concentrations, which were measured at 5 min intervals using the glucose oxidase method (2300 Stat; Yellow Springs Instruments, Yellow Springs, OH). A third cannula in the other antecubital fossa was dedicated to blood sampling for inflammatory markers.

On each study day, the arterialized blood glucose was stabilized initially at 4.5 mmol/l for 30 min and either maintained at that level (euglycemia) or lowered over 20 min to 2.5 mmol/l for 60 min (hypoglycemia), after which blood glucose was restored to 4.5 mmol/l. Subjects consumed a standardized meal after each study. Blood sample time points were: baseline, during the experimental session (+45 min), during recovery (+105 min), at +6 h, and at +24 h.

Flow cytometry

Whole blood samples were collected at the predetermined time points using D-Phenylalanyl-L-prolyl-L-arginine chloromethyl ketone, a selective thrombin inhibitor, as an anticoagulant. Samples (100 μl) of whole blood were immediately incubated with 10 μl of each monoclonal antibody (AbD Serotec, Kidlington, U.K.) for 30 min at room temperature, with subsequent red cell lysis by the addition of 1 ml of fluorescent-activated cell sorter (FACS) Lyse solution (Becton Dickinson, Oxford, U.K.). Flow cytometry using the FACS Calibur system (Becton Dickinson, Oxford, U.K.) was performed immediately after the experimental session to assess platelet-monocyte aggregation (CD14/CD42a) and CD40 expression on monocytes (CD14/CD40). Isotype controls were performed in addition to both mono- and dual-stain for each parameter assessed at each time point.

Soluble marker assays

Citrated plasma and serum samples were collected at the predetermined time points. These were separated immediately and frozen at –80°C until analysis for the soluble markers:

Von Willebrand factor (vWF) (enzyme-linked immunosorbent assay [ELISA]; coefficient of variation [CV] 7.3%), tissue plasminogen activator (tPA) antigen (Hyphen Biomed Zymutest; intra-assay CV 3.5%, inter-assay CV 4.4%), soluble CD40 ligand (sCD40L) (high sensitivity ELISA, Bender Medsystems; intra-assay CV 5.5%, inter-assay CV 7.2%), soluble P-selectin (ELISA, R&D Systems; intra-assay CV 5.1%, inter-assay CV 8.8%), IL-6 (High sensitivity ELISA, R&D Systems; intra-assay CV 5.9%, inter-assay CV 9.9%), and high sensitivity CRP (DRG Diagnostics; DRG Instruments, Marburg, Germany; intra-assay CV 4.2%, inter-assay CV 4.1%).

Catecholamine assays

Samples for epinephrine quantification were collected in EDTA tubes and immediately separated and frozen at –80°C until analysis by high-performance liquid chromatography and electrochemical detection (intra-assay CV 1.2%, inter-assay CV 3.9%).

Hypoglycemia symptom score

The Edinburgh Hypoglycemia Scale (15) was used to assess the symptoms experienced during each experimental session.

Statistical analyses

Results were analyzed using SPSS version 15.0 for Windows (SPSS, Chicago, IL). A general linear model (repeated-measures ANOVA) was used, with order of session (euglycemia-hypoglycemia or hypoglycemia-euglycemia) as a between-subjects factor, and condition (euglycemia or hypoglycemia) as a within-subjects factor, to compare hypoglycemia with euglycemia. Additional analysis using paired *t* tests was performed to assess the change in any given parameter from baseline. A *P* value <0.05 was considered to be significant. Results are reported as mean ± SD unless otherwise stated.

RESULTS— Hypoglycemia provoked a symptomatic response in all subjects with increased scores of autonomic ($P \leq 0.002$), neuroglycopenic ($P < 0.001$), and malaise ($P \leq 0.008$) symptoms compared with baseline. Comparison of baseline levels of inflammatory, endothelial and platelet markers in nondiabetic subjects and subjects with type 1 diabetes showed a significantly higher concentration of soluble P-selectin ($P = 0.01$) and of CD40 expression on monocytes ($P = 0.006$) in those with diabetes, demonstrating the chronic inflammatory response associated with diabetes.

Blood glucose

Target blood glucose concentrations were achieved (Fig. 1). In nondiabetic subjects, blood glucose concentrations were 2.58 ± 0.2 and 4.42 ± 0.5 mmol/l during hypoglycemia and euglycemia, respectively. In those with type 1 diabetes, blood glucose concentrations were 2.46 ± 0.22 and 4.53 ± 0.24 mmol/l, respectively. The blood glucose nadir was similar in both groups.

Counterregulatory response

Plasma epinephrine increased during hypoglycemia in participants with and with-




Figure 1—A: Blood glucose concentrations during hyperinsulinemic hypoglycemic and euglycemic clamp studies. B: Epinephrine responses to experimental procedures. i. nondiabetic subjects; ii. subjects with type 1 diabetes.

out type 1 diabetes ($P \leq 0.001$; Fig. 1). The epinephrine response occurred only during hypoglycemia and returned rapidly to baseline as anticipated (16).

Platelet activation

Platelet-monocyte aggregation. In nondiabetic subjects, platelet-monocyte aggregation appeared to rise, from a baseline level of $0.72 \pm 0.8\%$ to $3.09 \pm 8.1\%$ during hypoglycemia, with a peak of $3.49 \pm 10.4\%$ at 24 h (Fig. 2). Platelet-monocyte aggregation remained unchanged throughout euglycemia. The difference between conditions, and from baseline, did not achieve statistical significance.

In participants with diabetes, there was a late rise in platelet-monocyte aggregation after hypoglycemia at 24 h compared with baseline ($P = 0.03$).

Soluble P-selectin. Soluble plasma P-selectin concentrations increased after hypoglycemia in nondiabetic subjects, exhibiting a late response at 6 h ($P = 0.01$) and 24 h ($P = 0.02$; Fig. 2) but

decreasing during euglycemia ($P = 0.006$).

P-selectin also decreased during euglycemia in the diabetic group ($P = 0.04$), but did not change during hypoglycemia.

Endothelial markers

tPA. In nondiabetic subjects, plasma tPA concentrations increased during hypoglycemia, with a higher peak tPA concentration (12.55 ± 16.7 compared with 6.80 ± 7.9 ng/ml) (NS between conditions). Plasma tPA decreased significantly between baseline and test phase ($P = 0.004$) and recovery phase ($P = 0.006$), with a paradoxical rise between baseline and 24 h ($P = 0.06$) after euglycemia (Table 2). However, a diurnal variation in tPA concentration is recognized to occur, which may account for the decline observed during euglycemia (17). No significant differences occurred in the diabetic group (Table 2).

vWF. A trend toward a difference in plasma vWF concentrations was observed between hypoglycemia and euglycemia at

6 h in the nondiabetic subjects ($P = 0.07$) (Table 2).

Plasma vWF concentrations decreased between baseline and test phase ($P = 0.02$) and recovery phase ($P = 0.03$) after euglycemia in the participants with diabetes. No such decrement was observed during hypoglycemia (Table 2).

Inflammation

CD40 expression. CD40 expression on monocytes increased after hypoglycemia in nondiabetic subjects, from a baseline of $1.92 \pm 2.2\%$ to a maximum of $3.13 \pm 2.3\%$ at 24 h ($P = 0.009$). A significant difference between hypoglycemia and euglycemia conditions was present at 6 h ($P = 0.05$) and at 24 h ($P = 0.04$) (Table 2).

In participants with type 1 diabetes, monocyte CD40 expression increased from $3.69 \pm 3.4\%$ to $5.54 \pm 4.4\%$ during hypoglycemia ($P = 0.006$), compared with no change during euglycemia ($3.64 \pm 2.0\%$ to $3.65 \pm 1.8\%$, respectively; $P = \text{NS}$). The increment during




Figure 2—Platelet activation in response to experimental hypoglycemia and euglycemia. A: Platelet-monocyte aggregation. B: Soluble P-selectin. i. Nondiabetic subjects, ii. Subjects with type 1 diabetes.

hypoglycemia had dissipated by the time of the recovery phase and remained unchanged thereafter (Table 2).

sCD40L. In nondiabetic subjects, plasma sCD40L concentrations were higher during hypoglycemia than during euglycemia (2.80 ± 3.2 vs. 2.41 ± 2.8 ng/ml), with a trend toward significance ($P = 0.09$). A significant reduction in sCD40L concentration occurred during euglycemia between baseline and recovery phase ($P = 0.03$) (Table 2).

In those with diabetes, a significant difference was observed between the baseline levels on each study day: 3.36 ± 2.9 ng/ml on the hypoglycemia day compared with 2.86 ± 2.8 ng/ml on euglycemia day ($P = 0.03$), rendering subsequent measurements difficult to compare. A significant difference was again observed between the experimental condition levels, with a level of 3.41 ± 3.2 ng/ml during hypoglycemia and 2.85 ± 2.8 ng/ml during euglycemia ($P = 0.03$) (Table 2). Changes from baseline did not achieve significance.

IL-6. IL-6 levels rose in all experiments, maximally at 6 h, irrespective of

condition, with no clear differences identifiable in either group between the study conditions (Table 2).

hsCRP. Test phase hsCRP was higher in all subjects during hypoglycemia (1.81 ± 1.9 vs. 1.22 ± 1.9 ng/ml in nondiabetic participants [$P = 0.02$]; 2.72 ± 3.1 vs. 2.20 ± 2.9 ng/ml in subjects with diabetes [$P = \text{ns}$]) (Table 2). A significant difference was observed in the baseline concentrations in the nondiabetic participants ($P = 0.01$), frustrating interpretation of subsequent responses.

CONCLUSIONS— Previous studies have demonstrated that hypercoagulability, platelet and neutrophil activation, C-reactive protein, IL-6, and Endothelin-1 are upregulated after acute hypoglycemia (3,6–11), while a euglycemic insulin infusion (for at least 2 h) was shown to reduce inflammatory markers, consistent with an anti-inflammatory effect of insulin (18). The present study sought to replicate these effects, while investigating other underlying mechanisms of vascular disease, and tests were selected to investigate the effect of acute hypo-

glycemia on important cellular processes (platelet activation, endothelial dysfunction and inflammation) underlying the development of acute and chronic vascular complications in type 1 diabetes.

The present study showed that hypoglycemia generated a response in some of these markers, suggesting that hypoglycemia-induced metabolic stress may have adverse pathophysiological consequences while the euglycemic insulin infusion caused a potentially beneficial decrement in some parameters. However, the magnitude of most observed changes was small, and not all markers changed significantly.

The present study confirmed that platelet activation is promoted by hypoglycemia (8), with increments both in platelet-monocyte aggregation and soluble P-selectin. Conversely, P-selectin decreased during euglycemia. Endothelial function, using vWF and tPA Ag as surrogate markers, may have been disrupted, as shown by the increase in vWF after hypoglycemia in nondiabetic volunteers, but this change was not replicated in those with diabetes. However, a reduc-

Table 2—Endothelial function and inflammation in nondiabetic subjects and subjects with type 1 diabetes

	Euglycemia					Hypoglycemia				
	Baseline	Test	Recovery	+6 h	+24 h	Baseline	Test	Recovery	+6 h	+24 h
Nondiabetic subjects										
tPA (ng/ml)	7.37 ± 8.1	6.80 ± 7.9*	6.44 ± 7.5*	6.99 ± 9.3	8.51 ± 7.8†	10.96 ± 11.8	12.55 ± 16.7	9.10 ± 10.2*	9.83 ± 12.0	11.45 ± 11.6
vWF (IU/ml)	0.81 ± 0.3	0.76 ± 0.3	0.78 ± 0.2	0.80 ± 0.3	0.85 ± 0.3	0.82 ± 0.3	0.81 ± 0.3	0.89 ± 0.5	0.90 ± 0.3	0.89 ± 0.3
CD40 (%)	1.51 ± 1.4	2.23 ± 3.2†	2.40 ± 3.2†	0.84 ± 0.7	2.06 ± 1.0	1.92 ± 2.2	1.47 ± 1.1	1.55 ± 1.5	1.98 ± 2.4†	3.13 ± 2.3††
sCD40L (ng/ml)	2.68 ± 3.1	2.41 ± 2.8	2.40 ± 2.9*	2.63 ± 2.9	3.08 ± 3.3	2.88 ± 3.3	2.80 ± 3.2	2.55 ± 3.2	2.72 ± 3.3	2.79 ± 3.2
IL-6 (pg/ml)	0.86 ± 0.5	1.06 ± 1.2	1.05 ± 1.0	5.98 ± 4.6†	1.23 ± 0.9	0.72 ± 0.4	0.92 ± 0.5	1.62 ± 1.2†	4.37 ± 4.3†	1.00 ± 0.9
hsCRP (ng/ml)	1.04 ± 1.1	1.22 ± 1.9	1.18 ± 1.9	1.24 ± 1.6	1.31 ± 1.5	1.83 ± 1.5†	1.81 ± 1.9†	1.56 ± 1.3*	1.69 ± 1.2	1.90 ± 1.6
Subjects with type 1 diabetes										
tPA (ng/ml)	15.25 ± 30.2	17.70 ± 31.1	15.99 ± 27.5	22.13 ± 46.2	20.86 ± 34.8	18.12 ± 30.1	20.55 ± 36.1	17.69 ± 31.1	18.37 ± 32.6	22.98 ± 40.3
vWF (IU/ml)	0.91 ± 0.2	0.85 ± 0.2*	0.91 ± 0.3	0.85 ± 0.2*	0.99 ± 0.2	0.93 ± 0.2	0.95 ± 0.2	0.91 ± 0.2	0.90 ± 0.2	1.02 ± 0.2
CD40 (%)	3.64 ± 2.0	3.65 ± 1.8	4.14 ± 2.5	3.97 ± 2.3	4.35 ± 2.0	3.69 ± 3.4	5.54 ± 4.4†	3.36 ± 3.0	4.88 ± 2.4	4.70 ± 2.8
sCD40L (ng/ml)	2.86 ± 2.8	2.85 ± 2.8	2.84 ± 2.8	2.91 ± 2.9	3.25 ± 3.2†	3.36 ± 2.9†	3.41 ± 3.2†	3.10 ± 2.9*	3.05 ± 2.8*	3.44 ± 2.9
IL-6 (pg/ml)	0.69 ± 0.5	1.38 ± 1.9	1.58 ± 1.8	2.25 ± 2.8†	1.19 ± 1.2	1.21 ± 1.7	1.15 ± 1.5	1.76 ± 1.5	3.10 ± 4.9	1.96 ± 2.2
hsCRP (ng/ml)	2.52 ± 3.1	2.20 ± 2.9	2.32 ± 2.8	1.92 ± 1.8	3.40 ± 3.6	2.84 ± 3.2	2.72 ± 3.1	2.70 ± 3.2	2.89 ± 3.3	2.34 ± 2.8

Data are means ± SD. *Significant decrease from baseline ($P < 0.05$); †Significant increase from baseline ($P < 0.05$); ‡Significant difference between hypoglycemia and euglycemia ($P < 0.05$).

tion in vWF occurred after euglycemia in diabetic participants, which should confer vascular benefit. tPA Ag also appeared to increase in nondiabetic subjects during hypoglycemia, while declining during euglycemia, whereas no significant changes occurred in the diabetic group. Soluble markers of inflammation, sCD40L and hsCRP, were higher during hypoglycemia, with an elevation of hsCRP being observed in all subjects. Unfortunately, baseline differences in hsCRP in nondiabetic subjects, and in sCD40L in the diabetic subjects, frustrated interpretation of subsequent responses. sCD40L was apparently reduced during euglycemia in nondiabetic participants. Surprisingly, IL-6 increased in all experiments regardless of glycemic status, with a maximal response at 6 h. This response is inexplicable, and contrasts with a previous report (10). Monocyte CD40 expression also increased, suggesting promotion of the interaction of the CD40-CD40 ligand dyad (from the tumor necrosis factor receptor family), thus affecting another process in the pathway leading to atherosclerotic plaque rupture (19,20). This change occurred much earlier in the diabetic than the nondiabetic subjects, in whom the response was delayed, prolonged, and still present at 24 h. The persistence of these vascular changes for 24 h after the hypoglycemic stimulus, or their later emergence, suggests that the period of risk after hypoglycemia may be present long after blood glucose recovery.

For some markers, a positive trend after hypoglycemia was evident, without achieving statistical significance, or the only measurable difference between conditions was a beneficial effect associated with euglycemia. The sample size may have been insufficient to achieve significance, particularly as the magnitude of responses was small. It was not feasible to study a larger number of subjects using a procedure that is labor-intensive and costly. In a previous study, larger increments in inflammatory markers were observed during an insulin tolerance test, where hypoglycemia of <39 mg/dl (<2.2 mmol/l) was induced (21). The more rapid reduction to a lower blood glucose causing a greater hypoglycemic stimulus may have heightened the magnitude of the responses, compared with the more modest changes that occurred during a controlled glucose clamp (blood glucose 2.5 mmol/l [45 mg/dl]), as observed in the present study. A further limitation of the present study was the need to ex-

amine the experimental conditions on two separate days in a counterbalanced fashion. Because the baseline levels of many inflammatory markers can differ on separate days, as was observed with sCD40L and hsCRP, this biological variability hinders the interpretation and comparison of subsequent results. However, the present study design was necessary to allow comparison of the euglycemia and hypoglycemia conditions in individual subjects, as both time and insulin infusion per se may exert effects on biomarker levels. This study design cannot control for other day-to-day factors that could influence baseline levels of inflammatory markers. However, the effects of hypoglycemia could be evaluated, as each participant acted as their own control. This produces less variability than a comparison of results among individuals, as more inter-individual variation in inflammatory marker levels is present than intra-individual variation. In addition, it was possible to analyze each study separately, by examining changes in parameters from baseline on that particular day, enabling the detection of significant effects exerted by hypoglycemia compared with euglycemia. Baseline levels of all markers (except IL-6) were higher in the diabetic group (significant for P-selectin and CD40 expression). This could affect the magnitude of response induced by the experimental procedures. However, an analysis of the percentage change from baseline was consistent with the trends identified in the absolute results (shown as in the online appendix available at <http://care.diabetesjournals.org/cgi/content/full/dc10-0013/DC1>).

As anticipated, epinephrine secretion was stimulated by hypoglycemia. It is likely that hormonal changes underlie the activation and upregulation of the vascular biomarkers. Catecholamines promote platelet activation (22), while adrenoceptor blockade attenuates these effects (23,24). The participants with type 1 diabetes exhibited attenuated plasma epinephrine responses to hypoglycemia compared with the nondiabetic subjects, who were naive to such a hypoglycemic stimulus, this being consistent with the recognized decline in the magnitude of counterregulatory hormonal responses with increasing duration of type 1 diabetes (25). This attenuated epinephrine response may explain the lower responses of vascular biomarkers to hypoglycemia.

In summary, the effects of hypogly-

cemia on several vascular biomarkers that are implicated in the pathogenesis of vascular disease, would support the premise that acute hypoglycemia may be detrimental to an already diseased vasculature (2). Euglycemia may have a protective, anti-inflammatory effect. In the present study, the participants had no overt vascular disease and were unlikely to develop any demonstrable effects from a short period of exposure to hypoglycemia. However, in people with diabetes of long duration, who are likely to have underlying vascular disease, these responses may not be benign. The release of potent vasoactive substances could potentially aggravate chronic vasculopathy, and contribute to the precipitation of acute macrovascular events. This may aggravate established diabetic micro- and macrovascular disease in those who are exposed to recurrent hypoglycemia.

Acknowledgments—The cost of assays was supported by research grants from the Scottish Society of Physicians and the Edinburgh branch of Diabetes UK.

No potential conflicts of interest relevant to this article were reported.

References

- Hanssen KF, Dahl-Jørgensen K, Lauritzen T, Feldt-Rasmussen B, Brinchmann-Hansen O, Deckert T. Diabetic control and microvascular complications: the near-normoglycaemic experience. *Diabetologia* 1986;29:677–684
- Frier BM, Hilsted J. Does hypoglycaemia aggravate the complications of diabetes? *Lancet* 1985;2:1175–1177
- Wright RJ, Frier BM. Vascular disease and diabetes: is hypoglycaemia an aggravating factor? *Diabetes Metab Res Rev* 2008;24:353–363
- Action to Control Cardiovascular Risk in Diabetes Study Group, Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545–2559
- Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD, Zieve FJ, Marks J, Davis SN, Hayward R, Warren SR, Goldman S, McCarren M, Vitek ME, Henderson WG, Huang GD, for the VADT Investigators. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009;360:129–139
- Frier BM, Corral RJ, Davidson NM, Web-

- ber RG, Dewar A, French EB. Peripheral blood cell changes in response to acute hypoglycaemia in man. *Eur J Clin Invest* 1983;13:33–39
- Fisher BM, Quin JD, Rumley A, Lennie SE, Small M, MacCuish AC, Lowe GD. Effects of acute insulin-induced hypoglycaemia on haemostasis, fibrinolysis and haemorheology in insulin-dependent diabetic patients and control subjects. *Clin Sci* 1991;80:525–531
- Trovati M, Anfossi G, Cavalot F, Vitali S, Massucco P, Mularoni E, Schinco P, Tampone G, Emanuelli G. Studies on mechanisms involved in hypoglycemia-induced platelet activation. *Diabetes* 1986;35:818–825
- Galloway PJ, Thomson GA, Fisher BM, Semple CG. Insulin-induced hypoglycemia induces a rise in C-reactive protein (Letter). *Diabetes Care* 2000;23:861
- Dotson S, Freeman R, Failing HJ, Adler GK. Hypoglycemia increases serum interleukin-6 levels in healthy men and women. *Diabetes Care* 2008;31:1222–1223
- Wright RJ, Macleod KM, Perros P, Johnston N, Webb DJ, Frier BM. Plasma endothelin response to acute hypoglycaemia in adults with type 1 diabetes. *Diabet Med* 2007;24:1039–1042
- Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation* 2002;105:1135–1143
- De Fronzo R, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 1979;273:E214–E223
- Abumrad NN, Rabin D, Diamond MP, Lacy WW. Use of a heated superficial hand vein as an alternative site for the measurement of amino acid concentrations and for the study of glucose and alanine kinetics in man. *Metabolism* 1981;30:936–940
- Gold AE, MacLeod KM, Frier BM. Frequency of severe hypoglycemia in patients with type 1 diabetes with impaired awareness of hypoglycemia. *Diabetes Care* 1994;17:697–703
- Thompson CJ, Baylis PH. Endocrine changes during insulin-induced hypoglycaemia. In *Hypoglycaemia and Diabetes: Clinical and Physiological Aspects*. Frier BM, Fisher BM, Eds. Edward Arnold, London, U.K., 1993, p.116–131
- Rydzewski A, Urano T, Nagai N, Takada Y, Katoh-Oishi Y, Taminato T, Yoshimi T, Takada A. Diurnal variation in serum remnant-like lipoproteins, platelet aggregation and fibrinolysis in healthy volunteers. *Haemostasis* 1997;27:305–314
- Dandona P, Chaudhuri A, Ghanim H, Mohanty P. Insulin as an anti-inflammatory and antiatherogenic modulator. *J Am Coll Cardiol* 2009;53 (Suppl. 5):S14–S20
- Schönbeck U, Libby P. CD40 signaling and plaque instability. *Circ Res* 2001;89:1092–1103

20. Mach F, Schönbeck U, Libby P. CD40 signaling in vascular cells: a key role in atherosclerosis? *Atherosclerosis* 1998; 137(Suppl.):S89–S95
21. Razavi Nematollahi L, Kitabchi AE, Kitabchi AE, Stentz FB, Wan JY, Larijani BA, Tehrani MM, Gozashti MH, Omidfar K, Taheri E. Proinflammatory cytokines in response to insulin-induced hypoglycemic stress in healthy subjects. *Metabolism* 2009;58:443–448
22. Steel CM, French EB, Aitchison WR. Studies on adrenaline-induced leucocytosis in normal man. I. The role of the spleen and of the thoracic duct. *Br J Haematol* 1971;21:413–421
23. Fisher BM, Hepburn DA, Smith JG, Frier BM. The effect of alpha-adrenergic blockade on responses of peripheral blood cells to acute insulin-induced hypoglycaemia in humans. *Eur J Clin Invest* 1990; 20:51–55
24. Takeda H, Kishikawa H, Shinohara M, Miyata T, Suzaki K, Fukushima H, Ichinose K, Shichiri M. Effect of alpha 2-adrenoceptor antagonist on platelet activation during insulin-induced hypoglycaemia in type 2 (noninsulin-dependent) diabetes mellitus. *Diabetologia* 1988;31:657–663
25. Kerr D, Richardson T. Counterregulatory deficiencies in diabetes. In *Hypoglycaemia in Clinical Diabetes*. 2nd edition. Frier BM, Fisher M, Eds. John Wiley and Sons, Chichester, U.K., 2007, p. 121–140